

Tilburg University

Limiting distributions of moment- and quantile-based measures for skewness and kurtosis

Moors, J.J.A.; Coenen, V.M.J.; Heuts, R.M.J.

Publication date:
1993

[Link to publication in Tilburg University Research Portal](#)

Citation for published version (APA):

Moors, J. J. A., Coenen, V. M. J., & Heuts, R. M. J. (1993). *Limiting distributions of moment- and quantile-based measures for skewness and kurtosis*. (Research memorandum / Tilburg University, Department of Economics; Vol. FEW 620). Unknown Publisher.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.


- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

ECO
CBM
R
7626
1993
620

UNIVERSITEIT
BRABANT



Rm

**LIMITING DISTRIBUTIONS OF MOMENT-
AND QUANTILE-BASED MEASURES FOR
SKEWNESS AND KURTOSIS**

J.J.A. Moors, V.M.J. Coenen, R.M.J. Heuts

FEW 620

Communicated by Prof.dr. B.B. van der Genugten



K.U.B.
BIBLIOTHEEK
TILBURG

Limiting distributions of moment- and quantile-based measures for skewness and kurtosis

J.J.A. Moors* V.M.J. Coenen* R.M.J. Heuts *

Abstract

Consider a random variable with distribution function F and central moments μ_i . Well-known measures for skewness and kurtosis of this distribution are given by the moment-based pair

$$(\beta_1, \beta_2) = (\mu_3/\mu_2^{3/2}, \mu_4/\mu_2^2)$$

Another pair of measures is based on the quantiles E_i defined by $F(E_i) = i/8$ (for continuous F). This quantile-based pair (S, T) is defined by

$$S = (E_6 - 2E_4 + E_2)/(E_6 - E_2)$$

$$T = (E_7 - E_5 + E_3 - E_1)/(E_6 - E_2)$$

Let (\hat{b}_1, \hat{b}_2) and (\hat{s}, \hat{t}) denote the natural estimators for (β_1, β_2) and (S, T) , respectively, based on a random sample of size n . In this paper, for both pairs of random variables the limiting distributions for $n \rightarrow \infty$ are derived. The results are checked by means of simulations.

Two possible applications of these limiting distributions are discussed briefly: modelling and tests of normality. Here, modelling (or curve-fitting) means: describing a given empirical distribution by means of a simple theoretical probability distribution. This modelling can be based either on the pair (\hat{b}_1, \hat{b}_2) or on the pair (\hat{s}, \hat{t}) ; the asymptotic efficiency of the two methods depends on the limiting distributions a.o. A test of normality based on (\hat{b}_1, \hat{b}_2) is well-known; the pair (\hat{s}, \hat{t}) presents an alternative test statistic. The asymptotic power of both tests can be found from the limiting distributions.

*Tilburg University, P.O. Box 90153, 5000 LE Tilburg, Netherlands

1 The limiting distribution of $(\underline{b}_1, \underline{b}_2)$

Consider a random variable \underline{x} with moments $\alpha_i = E\{\underline{x}^i\}$; assume $\alpha_8 < \infty$. From

$$Cov(\underline{x}^i, \underline{x}^j) = \alpha_{i+j} - \alpha_i \alpha_j \quad (i, j = 1, 2, 3, 4)$$

the covariance-matrix Σ_4 of $(\underline{x}, \underline{x}^2, \underline{x}^3, \underline{x}^4)$ follows:

$$\Sigma_4 = \begin{pmatrix} \alpha_2 - \alpha_1^2 & \alpha_3 - \alpha_1 \alpha_2 & \alpha_4 - \alpha_1 \alpha_3 & \alpha_5 - \alpha_1 \alpha_4 \\ \alpha_3 - \alpha_1 \alpha_2 & \alpha_4 - \alpha_2^2 & \alpha_5 - \alpha_2 \alpha_3 & \alpha_6 - \alpha_2 \alpha_4 \\ \alpha_4 - \alpha_1 \alpha_3 & \alpha_5 - \alpha_2 \alpha_3 & \alpha_6 - \alpha_3^2 & \alpha_7 - \alpha_3 \alpha_4 \\ \alpha_5 - \alpha_1 \alpha_4 & \alpha_6 - \alpha_2 \alpha_4 & \alpha_7 - \alpha_3 \alpha_4 & \alpha_8 - \alpha_4^2 \end{pmatrix} \quad (1.1)$$

For a random sample $(\underline{x}_1, \underline{x}_2, \dots, \underline{x}_n)$ the sample moments \underline{a}_i are defined by

$$\underline{a}_i = \frac{1}{n} \sum_{t=1}^n \underline{x}_t^i \quad (i = 1, 2, 3, 4)$$

Then the multivariate version of the familiar Lindeberg-Lévy theorem gives

$$\sqrt{n} \begin{pmatrix} \underline{a}_1 - \alpha_1 \\ \underline{a}_2 - \alpha_2 \\ \underline{a}_3 - \alpha_3 \\ \underline{a}_4 - \alpha_4 \end{pmatrix} \xrightarrow{L} \underline{y} \sim N_4(0, \Sigma_4) \quad (1.2)$$

where \xrightarrow{L} denotes convergence in distribution. See SERFLING (1980), p. 68.

To find the limiting distribution of the sample statistics

$$\begin{pmatrix} \underline{b}_1 \\ \underline{b}_2 \end{pmatrix} = \begin{pmatrix} \frac{1}{n} \sum_t (\underline{x}_t - \bar{\underline{x}})^3 / \left[\frac{1}{n} \sum_t (\underline{x}_t - \bar{\underline{x}})^2 \right]^{3/2} \\ \frac{1}{n} \sum_t (\underline{x}_t - \bar{\underline{x}})^4 / \left[\frac{1}{n} \sum_t (\underline{x}_t - \bar{\underline{x}})^2 \right]^2 \end{pmatrix}$$

for skewness and kurtosis the following well-known result will be used.

Theorem 1. Let $\underline{y}_n, \underline{y}, c \in \mathbf{R}^k$ and $\varphi : \mathbf{R}^k \rightarrow \mathbf{R}^m$ with φ totally differentiable in c . If

$$\sqrt{n}(\underline{y}_n - c) \xrightarrow{L} \underline{y}$$

holds, then

$$\sqrt{n}[\varphi(\underline{y}_n) - \varphi(c)] \xrightarrow{L} \varphi'(c)\underline{y}$$

holds as well. \square

Note that $\varphi'(c)$ is a $m \times k$ -matrix of partial derivatives. Choose

$$\begin{aligned}\underline{y}_n &= (\underline{a}_1, \underline{a}_2, \underline{a}_3, \underline{a}_4)^\top \\ c &= (\alpha_1, \alpha_2, \alpha_3, \alpha_4)^\top\end{aligned}$$

$$\varphi(y_1, y_2, y_3, y_4) = \begin{pmatrix} [y_3 - 3y_2y_1 + 2y_1^3]/[y_2 - y_1^2]^{3/2} \\ [y_4 - 4y_3y_1 + 6y_2y_1^2 - 3y_1^4]/[y_2 - y_1^2]^2 \end{pmatrix}$$

Then (1.2) may be rewritten as

$$\sqrt{n}(\underline{y}_n - c) \xrightarrow{L} \underline{y} \sim N_4(0, \Sigma_4)$$

Since φ is totally differentiable at c and

$$\begin{aligned}\varphi(\underline{y}_n) &= \varphi(\underline{a}_1, \underline{a}_2, \underline{a}_3, \underline{a}_4) = (\underline{b}_1, \underline{b}_2)^\top \\ \varphi(c) &= \varphi(\alpha_1, \alpha_2, \alpha_3, \alpha_4) = (\beta_1, \beta_2)^\top\end{aligned}$$

Theorem 1 implies

$$\sqrt{n} \begin{pmatrix} \underline{b}_1 - \beta_1 \\ \underline{b}_2 - \beta_2 \end{pmatrix} \xrightarrow{L} \varphi'(c)\underline{y} \sim N_2(0, \Sigma^M) \quad (1.3)$$

where $\Sigma^M = \varphi'(c) \Sigma_4 [\varphi'(c)]^\top$; the superscript M stands for 'moments'.

The explicit calculation of Σ^M can be simplified greatly by observing that this covariance-matrix is location-invariant. Hence, the calculation can be carried out for the special case $\mu_1 = 0$, implying $\alpha_i = \mu_i$ for all i . This gives

$$\Sigma^M = \varphi'(c)\Sigma_4[\varphi'(c)]^\top = AVA^\top$$

with

$$A = \begin{pmatrix} -3/\mu_2^{1/2} & -3\mu_3/2\mu_2^{5/2} & \mu_2 & 0 \\ -4\mu_3/\mu_2^2 & -2\mu_4/\mu_2^3 & 0 & 1/\mu_2^2 \end{pmatrix}$$

$$V = \begin{pmatrix} \mu_2 & \mu_3 & \mu_4 & \mu_5 \\ \mu_3 & \mu_4 - \mu_2^2 & \mu_5 - \mu_2\mu_3 & \mu_6 - \mu_2\mu_4 \\ \mu_4 & \mu_5 - \mu_2\mu_3 & \mu_6 - \mu_2^2 & \mu_7 - \mu_3\mu_4 \\ \mu_5 & \mu_6 - \mu_2\mu_4 & \mu_7 - \mu_3\mu_4 & \mu_8 - \mu_4^2 \end{pmatrix}$$

Then the elements Σ_{ij}^M of the covariance-matrix Σ^M are easily found to be

$$\Sigma_{11}^M = \frac{\mu_6}{\mu_2^3} - 3\beta_1 \frac{\mu_5}{\mu_2^{5/2}} + \beta_1^2(35 + 9\beta_2)/4 - 6\beta_2 + 9$$

$$\Sigma_{12}^M = \Sigma_{21}^M = \frac{\mu_7}{\mu_2^{7/2}} - \frac{3}{2}\beta_1 \frac{\mu_6}{\mu_2^3} - (2\beta_2 + 3) \frac{\mu_5}{\mu_2^{5/2}} + 3\beta_1(\beta_2^2 + \beta_2/2 + 2\beta_1^2 + 4)$$

$$\Sigma_{22}^M = \frac{\mu_8}{\mu_2^4} - 4\beta_2 \frac{\mu_6}{\mu_2^3} - 8\beta_1 \frac{\mu_5}{\mu_2^{5/2}} + \beta_2^2(4\beta_2 - 1) + 16\beta_1^2(\beta_2 + 1)$$

The expressions for Σ_{11}^M and Σ_{22}^M can be found in STUART & ORD (1987), p. 344; the derivation was quite different, however. The formula for the limiting covariance of \underline{b}_1 and \underline{b}_2 seems to be new.

For the standardnormal and the standardexponential distribution Σ^M was calculated directly as well, with the following results:

$$N(0, 1) \Rightarrow \Sigma^M = \begin{pmatrix} 6 & 0 \\ 0 & 24 \end{pmatrix}$$

$$Ne(1) \Rightarrow \Sigma^M = \begin{pmatrix} 72 & 720 \\ 720 & 8064 \end{pmatrix}$$

These outcomes are in agreement with the general formulae.

2 The limiting distribution of $(\underline{s}, \underline{t})$

For any random variable \underline{x} a p -quantile Q_p is defined by

$$P(\underline{x} < Q_p) \leq p, \quad P(\underline{x} \geq Q_p) \leq 1 - p$$

for $0 < p < 1$. If the distribution function F of \underline{x} is differentiable in Q_p with $F'(Q_p) > 0$, this definition can be simplified to

$$F(Q_p) = p$$

Now, consider m given numbers p_i satisfying

$$0 < p_1 < p_2 < \dots < p_m < 1$$

and denote the corresponding p_i -quantiles by Q_i . Let \underline{q}_i denote the sample p_i -quantile ($i = 1, 2, \dots, m$) for a random sample $(\underline{x}_1, \underline{x}_2, \dots, \underline{x}_n)$ from distribution function F . Then the following limit theorem holds; see SERFLING (1980), p. 80.

Theorem 2. If F is differentiable in Q_i with derivative $f(Q_i) = F'(Q_i) > 0$ ($i = 1, 2, \dots, m$), then

$$\sqrt{n} \begin{pmatrix} \underline{q}_1 - Q_1 \\ \underline{q}_2 - Q_2 \\ \vdots \\ \underline{q}_m - Q_m \end{pmatrix} \xrightarrow{L} \underline{y} \sim N_m(0, \Sigma)$$

where $\Sigma_{ij} = p_i(1 - p_j)/[f(Q_i)f(Q_j)]$ for $i \leq j$ and $\Sigma_{ij} = \Sigma_{ji}$. \square

The seven quantiles corresponding with $p_i = i/8$ are called octiles and will be denoted by E_i , $i = 1, 2, \dots, 7$. Alternative measures for skewness and kurtosis exist, based on these octiles:

$$\begin{cases} S = (E_6 - 2E_4 + E_2)/(E_6 - E_2) \\ T = (E_7 - E_5 + E_3 - E_1)/(E_6 - E_2) \end{cases} \quad (2.1)$$

Of course, the quantile-measure S is the well-known quantile-coefficient of skewness; for the properties of T see MOORS (1988) or WAGEMAKERS et al. (1992).

Now the limiting distribution of the seven sample octiles \underline{e}_i is an immediate consequence of Theorem 2.

Corollary 1. If F is differentiable in E_i with derivative $f(E_i) = F'(E_i) > 0$ ($i = 1, 2, \dots, 7$), then

$$\sqrt{n} \begin{pmatrix} \underline{e}_1 - E_1 \\ \underline{e}_2 - E_2 \\ \vdots \\ \underline{e}_7 - E_7 \end{pmatrix} \xrightarrow{L} \underline{y} \sim N_7(0, \Sigma_7)$$

where

$$\Sigma_7 = \frac{1}{64} \begin{pmatrix} 7c_1^2 & 6c_1c_2 & 5c_1c_3 & 4c_1c_4 & 3c_1c_5 & 2c_1c_6 & c_1c_7 \\ 6c_1c_2 & 12c_2^2 & 10c_2c_3 & 8c_2c_4 & 6c_2c_5 & 4c_2c_6 & 2c_2c_7 \\ 5c_1c_3 & 10c_2c_3 & 15c_3^2 & 12c_3c_4 & 9c_3c_5 & 6c_3c_6 & 3c_3c_7 \\ 4c_1c_4 & 8c_2c_4 & 12c_3c_4 & 16c_4^2 & 12c_4c_5 & 8c_4c_6 & 4c_4c_7 \\ 3c_1c_5 & 6c_2c_5 & 9c_3c_5 & 12c_4c_5 & 15c_5^2 & 10c_5c_6 & 5c_5c_7 \\ 2c_1c_6 & 4c_2c_6 & 6c_3c_6 & 8c_4c_6 & 10c_5c_6 & 12c_6^2 & 6c_6c_7 \\ c_1c_7 & 2c_2c_7 & 3c_3c_7 & 4c_4c_7 & 5c_5c_7 & 6c_6c_7 & 7c_7^2 \end{pmatrix}$$

with $c_i = 1/f(E_i)$. \square

The natural estimators for S and T are the sample statistics

$$\begin{cases} \underline{s} = (\underline{e}_6 - 2\underline{e}_4 + \underline{e}_2)/(\underline{e}_6 - \underline{e}_2) \\ \underline{t} = (\underline{e}_7 - \underline{e}_5 + \underline{e}_3 - \underline{e}_1)/(\underline{e}_6 - \underline{e}_2) \end{cases} \quad (2.2)$$

Their limiting distribution can be found with the help of Theorem 1. By taking

$$\begin{aligned} \underline{y}_n &= (\underline{e}_1, \underline{e}_2, \dots, \underline{e}_7)^\top \\ \underline{c} &= (E_1, E_2, \dots, E_7)^\top \end{aligned}$$

$$\varphi(y_1, y_2, \dots, y_7) = \begin{pmatrix} (y_6 - 2y_4 + y_2)/(y_6 - y_2) \\ (y_7 - y_5 + y_3 - y_1)/(y_6 - y_2) \end{pmatrix}$$

the result of Corollary 1 can be written as

$$\sqrt{n}(\underline{y}_n - c) \xrightarrow{L} \underline{y} \sim N_7(0, \Sigma_7)$$

Then Theorem 1 gives

$$\sqrt{n} \begin{pmatrix} \underline{s} - S \\ \underline{t} - T \end{pmatrix} \xrightarrow{L} \varphi'(c)\underline{y} \sim N_2(0, \Sigma^Q) \quad (2.3)$$

where $\Sigma^Q = \varphi'(c) \Sigma_7 [\varphi'(c)]^T$; the Q stands for 'quantiles'. With the notation

$$D_6 = E_6 - E_4, \quad D_4 = E_4 - E_2, \quad D = E_6 - E_2$$

the elements Σ_{ij}^Q of Σ^Q are found to be given by

$$\begin{aligned} 4D^4 \Sigma_{11}^Q &= (c_2 D_6 + c_6 D_4)^2 + 2(c_2 D_6 - c_4 D)^2 + 2(c_6 D_4 - c_4 D)^2 \\ 16D^3 \Sigma_{12}^Q &= c_2 D_6 (-3c_1 + 6c_2 T + 5c_3 - 3c_5 - 2c_6 T + c_7) + \\ &\quad + c_4 (2c_1 - 4c_2 T - 6c_3 + 6c_5 + 4c_6 T - 2c_7) + \\ &\quad + c_6 D_4 (-c_1 + 2c_2 T + 3c_3 - 5c_5 - 6c_6 T + 3c_7) \\ \Sigma_{21}^Q &= \Sigma_{12}^Q \\ 64D^2 \Sigma_{22}^Q &= 6(c_1 - c_2 T)^2 + 5(c_1 - c_3)^2 - 3(c_1 - c_5)^2 - 2(c_1 - c_6 T)^2 + \\ &\quad + (c_1 - c_7)^2 + 10(c_2 T + c_3)^2 - 6(c_2 T + c_5)^2 + 4(c_2 T - c_6 T)^2 + \\ &\quad - 2(c_2 T - c_7)^2 + 9(c_3 - c_5)^2 - 6(c_3 + c_6 T)^2 - 3(c_3 - c_7)^2 + \\ &\quad + 10(c_5 + c_6 T)^2 + 5(c_5 - c_7)^2 + 6(c_6 T - c_7)^2 \end{aligned}$$

The results for the two special distributions are:

$$\begin{aligned} N(0, 1) &\Rightarrow \Sigma^Q = \begin{pmatrix} 1.839 & 0 \\ 0 & 3.153 \end{pmatrix} \\ Ne(1) &\Rightarrow \Sigma^Q = \begin{pmatrix} 1.782 & -0.151 \\ -0.151 & 5.104 \end{pmatrix} \end{aligned}$$

3 Simulations

As a final check of the formulae for the covariance-matrices derived before, as well as to investigate how accurate the limiting distributions are for finite n , rather extensive simulation studies were undertaken. The core of these studies consisted of the following four steps.

1. Choose a certain probability distribution and calculate the pairs of measures (β_1, β_2) and (S, T) .
2. Simulate from this distribution a random sample of size n and calculate the pairs of statistics (b_1, b_2) and (s, t) .
3. Repeat step 2 k times, independently.
4. Calculate the means and the covariance-matrix for the k pairs $\sqrt{n}(b_1 - \beta_1, b_2 - \beta_2)$, as well as for the k pairs $\sqrt{n}(s - S, t - T)$.

In this procedure, the following choices were made.

- (i) Table 1 shows the distributions used in step 1.

Table 1. Distributions used in simulation studies.

type	parameters (p, q)			
$N(\mu, \sigma^2)$	(0,1)			
$B(p, q)$	(0.5,1)	(1,2)	(1,4)	(2,4)
$B2(p, q)$	(3,15)	(4,28)	(5,15)	
$\Gamma(p, q)$	(1,1)	(1,5)		
$I\Gamma(p, q)$	(1,9)	(1,12)		

Here, $B(p, q)$ indicates the beta-distribution with density

$$f(x) = x^{p-1}(1-x)^{q-1}/B(p, q), \quad 0 < x < 1; \quad p, q > 0$$

while $B2(p, q)$ is the beta-distribution of the second kind with density

$$f(x) = x^{p-1}/(1+x)^{p+q}/B(p, q), \quad x > 0; \quad p, q > 0$$

$\Gamma(p, q)$ denotes the gamma-distribution with density

$$f(x) = p^q x^{q-1} e^{-px} / \Gamma(q), \quad x > 0; \quad p, q > 0$$

and $I\Gamma(p, q)$ is the inverse gamma-distribution with density

$$f(x) = p^q x^{-q-1} e^{-p/x} / \Gamma(q) \quad x > 0; \quad p, q > 0$$

Note that all distributions in Table 1 belong to the Pearson-system of distributions; the reason for this is given in Section 4. Table 2 presents the values for the pairs (β_1, β_2) and (S, T) for the distributions selected in Table 1.

Table 2. Moments- and quantile-based measures
for skewness and kurtosis.

Distributions	β_1	β_2	S	T
$N(0, 1)$	0	3	0	1.233
$B(0.5, 1)$	0.639	2.143	0.25	1
$B(1, 2)$	0.566	2.4	0.132	1.103
$B(1, 4)$	1.050	3.696	0.197	1.190
$B(2, 4)$	0.468	2.625	0.078	1.167
$B2(3, 15)$	1.683	8.180	0.171	1.289
$B2(4, 28)$	1.282	5.819	0.139	1.266
$B2(5, 15)$	1.480	7.128	0.147	1.281
$\Gamma(1, 1)$	2	9	0.262	1.306
$\Gamma(1, 5)$	0.894	4.2	0.104	1.243
$I\Gamma(1, 9)$	1.764	9.8	0.151	1.302
$I\Gamma(1, 12)$	1.405	7.083	0.131	1.284

(If less than three decimals are shown, the value is exact.)

(ii) In step 2 the following sample sizes n were considered:

50, 100, 200, 2000

(ii) In step 3 two values were taken for the number of iterations k :

20, 200

Tables 3 and 4 show the detailed results for the standard normal distribution; the last lines for $n = \infty$ are obtained from the theoretical results Σ^M and Σ^Q in Section 1 and 2, respectively.

Table 3. Simulations and limiting values; $(\underline{b}_1, \underline{b}_2)$, $N(0, 1)$.

k	n	$\sqrt{n}E(\underline{b}_1 - \beta_1)$	$\sqrt{n}E(\underline{b}_2 - \beta_2)$	$nV(\underline{b}_1)$	$nV(\underline{b}_2)$	$n \text{Cov}(\underline{b}_1, \underline{b}_2)$
20	50	0.059	0.939	6.527	41.716	9.436
	100	-0.714	-1.596	5.067	19.704	-3.059
	200	-0.295	-0.968	4.889	33.844	-5.389
	2000	0.179	0.421	4.121	18.570	-1.890
200	50	-0.240	-0.893	5.068	15.099	-0.068
	100	0.095	-0.840	4.683	17.781	-0.035
	200	-0.013	-0.461	5.453	21.181	1.185
	2000	0.296	0.570	5.180	23.137	0.283
∞		0	0	6	24	0

Table 4. Simulations and limiting values; $(\underline{s}, \underline{t})$, $N(0, 1)$.

k	n	$\sqrt{n}E(\underline{s} - S)$	$\sqrt{n}E(\underline{t} - T)$	$nV(\underline{s})$	$nV(\underline{t})$	$n \text{Cov}(\underline{s}, \underline{t})$
20	50	-0.395	0.017	1.556	2.538	0.146
	100	0.548	0.391	1.757	4.341	0.402
	200	0.393	0.159	0.964	1.259	0.010
	2000	0.845	-0.170	1.532	1.969	0.330
200	50	-0.224	-0.149	1.887	3.940	0.256
	100	0.103	0.395	1.745	3.378	-0.034
	200	-0.039	0.395	1.931	3.487	-0.026
	2000	-0.048	0.654	1.524	3.815	-0.283
∞		0	0	1.839	3.153	0

As was to be expected, the simulated outcomes are closer to the limiting values when the sample size n and/or the number of iterations k increase.

To save space, only the simulated outcomes for the values $k = 200$ and $n = 2000$ are reported here (Tables 5 and 6) for the other distributions of Table 1. Fuller details can

be found in COENEN (1992).

Table 5. Simulations for $k = 200$ and $n = 2000$ ('sim') and limiting values ('lim'); (b_1, b_2) .

Distribution		$\sqrt{n}E(b_1 - \beta_1)$	$\sqrt{n}E(b_2 - \beta_2)$	$nV(b_1)$	$nV(b_2)$	$n \text{Cov}(b_1, b_2)$
$B(0.5, 1)$	sim	0.034	0.137	2.752	8.582	4.429
	lim	0	0	2.735	8.144	4.313
$B(1, 2)$	sim	0.021	0.051	2.601	9.418	4.225
	lim	0	0	2.499	8.887	3.927
$B(1, 4)$	sim	-0.186	-0.526	4.492	57.668	14.709
	lim	0	0	5.125	67.962	17.213
$B(2, 4)$	sim	-0.072	-0.079	2.896	12.315	4.667
	lim	0	0	2.999	12.979	4.789
$B2(3, 15)$	sim	0.288	6.858	139.41	24811	1750
	lim	0	0	119.35	22747	1488
$B2(4, 28)$	sim	0.364	4.515	49.038	3200	369.44
	lim	0	0	39.885	2944	315.03
$B2(5, 15)$	sim	-0.260	-2.777	81.860	8564	793.32
	lim	0	0	88.267	13493	982.04
$\Gamma(1, 1)$	sim	-0.274	-2.702	63.389	6481	615.68
	lim	0	0	72	8064	720
$\Gamma(1, 5)$	sim	0.103	-0.012	12.916	330.18	59.739
	lim	0	0	14.400	410.11	69.551
$I\Gamma(1, 9)$	sim	-0.221	-4.742	327.13	79197	4864
	lim	0	0	386.84	460077	9488
$I\Gamma(1, 12)$	sim	-0.659	-7.124	104.62	1891	1052
	lim	0	0	109.74	23002	1381

Table 6. Simulations for $k = 200$ and $n = 2000$ ('sim') and limiting values ('lim'); ($\underline{s}, \underline{t}$).

Distribution		$\sqrt{n}E(\underline{s} - S)$	$\sqrt{n}E(\underline{t} - T)$	$nV(\underline{s})$	$nV(\underline{t})$	$n \text{Cov}(\underline{s}, \underline{t})$
$B(0.5, 1)$	sim	-0.022	-0.015	1.956	2.662	0.061
	lim	0	0	2.092	2.623	0.094
$B(1, 2)$	sim	-0.083	-0.054	1.774	2.733	-0.135
	lim	0	0	1.900	2.648	-0.049
$B(1, 4)$	sim	0.097	0.020	1.802	3.744	-0.261
	lim	0	0	1.725	3.086	-0.136
$B(2, 4)$	sim	-0.058	0.101	1.938	3.190	0.228
	lim	0	0	1.854	2.847	-0.045
$B2(3, 15)$	sim	0.106	0.109	1.646	4.562	0.107
	lim	0	0	1.806	4.198	-0.115
$B2(4, 28)$	sim	-0.035	-0.015	1.589	4.343	-0.193
	lim	0	0	1.817	3.796	-0.093
$B2(5, 15)$	sim	0.038	0.316	1.580	3.424	-0.215
	lim	0	0	1.817	3.986	-0.101
$\Gamma(1, 1)$	sim	-0.050	-0.061	1.874	5.098	-0.340
	lim	0	0	1.783	5.104	-0.151
$\Gamma(1, 5)$	sim	0.070	0.099	1.967	3.393	0.096
	lim	0	0	1.830	3.445	-0.070
$I\Gamma(1, 9)$	sim	0.012	-0.161	1.663	4.236	-0.209
	lim	0	0	1.820	4.230	-0.110
$I\Gamma(1, 12)$	sim	0.088	0.030	1.408	4.307	-0.187
	lim	0	0	1.819	3.913	-0.091

The overall agreement is good, with the possible exception of the $V(\underline{b}_2)$ -values for the inverse-gamma-distributions. Note however, that these distributions have very high β_2 -values; probably, the number of replications was too small to obtain a good agreement.

4 Application in modelling


Modelling an empirical frequency distribution by means of a simple theoretical probability distribution is an interesting issue in applied statistics. One of the advantages is that the analytical expression for the density may lead to explicit cost formulae, e.g. in inventory problems. A reasonable approach to this modelling problem is to demand that measures for location, dispersion, skewness and kurtosis for the two distributions coincide. Traditionally, the moment-based quartet of measures $(\mu, \mu_2, \beta_1, \beta_2)$ was used hereby. An alternative is the quantile-based quartet (Q, R, S, T) , with the median Q and the half interquantile range R defined as

$$Q = E_4, \quad R = (E_6 - E_2)/2$$

So, this last quartet can be found from the seven octiles.

Of course, the values of such a foursome of measures do not uniquely determine a probability distribution in general. They do, however, within specific systems of distributions like Pearson's or Johnson's. Both systems have the property that any (location-scale-free) distribution within the system has a different pair of values for the measures (β_1, β_2) . Similarly, there is a one-one-correspondence between these (location-scale-free) distributions and the pairs (S, T) . Figure 1 shows these one-one-relations for the Pearson system; the (symmetric) half-planes with negative β_1 - and S -values have been omitted.

Figure 1. (β_1, β_2) - and (S, T) -plane for the Pearson system.



The Roman capitals indicate different types of distributions; e.g. I corresponds with the beta-distributions and III with the gamma distributions. N is the class of all normal distributions. See WAGEMAKERS (1992) for details.

Now, if for an empirical frequency distribution the values of the four moment-measures have been calculated, there is (at most) one distribution within the Pearson system with the same values for the quartet $(\mu, \mu_2, \beta_1, \beta_2)$. This unique distribution can be taken as a model for the empirical one. In the same way, the (unique) distribution within Pearson's system, having the same values of the quantile-based measures (Q, R, S, T) as the empirical distribution, presents another model.

The important question now is which of these two models is to be preferred: does the (classical) approach using $(\mu, \mu_2, \beta_1, \beta_2)$ lead to a better-fitting model than the quantile-based foursome (Q, R, S, T) does? The limiting distributions derived in this paper present a first step in answering this question: comparison of Tables 5 and 6 immediately shows that – for n large – $(\underline{s}, \underline{t})$ has (much) smaller variances than $(\underline{b}_1, \underline{b}_2)$. This implies that with the same size (S, T) can be estimated (much) more accurately than (β_1, β_2) .

However, comparing the accuracy of the two modelling methods is more complicated. Let $D(p, q)$ be one of the distributions in Pearson's system, D indicating the type; a random sample is drawn, leading to the pairs $(\underline{b}_1, \underline{b}_2)$ and $(\underline{s}, \underline{t})$. They lead to unique distributions, say

$$\underline{D}_M(\underline{p}_M, \underline{q}_M), \quad \underline{D}_Q(\underline{p}_Q, \underline{q}_Q)$$

respectively. The random variables $\underline{p}_M, \underline{p}_Q$ and $\underline{q}_M, \underline{q}_Q$ can be seen as estimators for p and q , respectively. Note, however, that the types \underline{D}_M and \underline{D}_Q are not necessarily identical to the sampled type D !

If $\underline{D}_M = \underline{D}_Q = D$ with probability 1, the accuracy of the modelling processes would only depend on the efficiency of the estimators $(\underline{p}_M, \underline{q}_M)$ and $(\underline{p}_Q, \underline{q}_Q)$, hence on the matrices of mean squares errors, S_M and S_Q , say. For the moment method, S_M is defined as

$$S_M = E[(\underline{p}_M - p, \underline{q}_M - q)(\underline{p}_M - p, \underline{q}_M - q)^T]$$

while a similar expression holds for S_Q . This method breaks down however, since in practice $P(\underline{D}_M = \underline{D}_Q = D_Q) = 1$ will never hold.

Some small-scale experiments were performed on this subject. From $D(p, q) = B(0.5, 1)$ twenty random samples of size $n = 50$ were drawn. In all cases but one the sample pairs (b_1, b_2) and (s, t) lead back to beta distributions – hence $D_M = D_Q = D$, the exception being a sample with $(s, t) = (0.241, 1.571)$, corresponding with a Pearson distribution of type IV. From the nineteen remaining samples we calculated the estimates \hat{S}_M and \hat{S}_Q , where e.g.

$$\hat{S}_M = \frac{1}{19} \sum_{i=1}^{19} (p_{Mi} - 0.5, q_{Mi} - 1)(p_{Mi} - 0.5, q_{Mi} - 1)^T$$

and i denotes the different sample. The results were

$$\hat{S}_M = \begin{pmatrix} 0.0296 & 0.0176 \\ 0.0176 & 0.0520 \end{pmatrix}, \quad \hat{S}_Q = \begin{pmatrix} 0.0202 & 0.0337 \\ 0.0337 & 0.1090 \end{pmatrix}$$

Since $\hat{S}_M - \hat{S}_Q$ is indefinite, no conclusion can be drawn about which modelling process is better - even if the problem of identifying the right **type** of distribution is neglected.

Better measures for the accuracy of the modelling methods can be obtained by comparing the distribution functions of the model and the sampled distributions. Denoting these by F_M (for the moment-based model) and F , respectively, useful measures of fit are the Kolmogorov-Smirnov-distance

$$\max_x |F_M(x) - F(x)|$$

or the Cramer-Von Mises statistic

$$\int_0^1 [F_M(x) - F(x)]^2 dF(x)$$

We plan to do some future research in this direction.

5 Application in test for normality

There exist quite a few tests for normality; SHAPIRO et al. (1968) discuss and compare nine. Among them are two tests using as test statistic \underline{b}_1 – in our notation – and \underline{b}_2 separately. These test statistics can be used in combination as well. D'AGOSTINO et al.

(1973) were the first to discuss a test based on the simultaneous distribution of $(\underline{b}_1, \underline{b}_2)$. The original testing problem:

- H_o : the sample stems from some normal distribution,
 H_a : the sample stems from any other distribution within the
 Pearson system

is in fact replaced by

$$H'_o : (\beta_1, \beta_2) = (0, 3), \quad H'_a : (\beta_1, \beta_2) \neq (0, 3)$$

The limiting distribution of $(\underline{b}_1, \underline{b}_2)$ obtained in Section 1 implies that for large n

$$\underline{b}_1 \approx N(0, 6/n), \quad \underline{b}_2 \approx N(3, 24/n), \quad \underline{b}_1 \text{ and } \underline{b}_2 \text{ independent}$$

holds under the null hypotheses. Here \approx means 'approximately distributed as'. Hence, under H_o

$$\underline{b} = n[\underline{b}_1^2/6 + (\underline{b}_2 - 3)^2/24] \approx \chi^2_2 \quad (5.1)$$

holds, so that \underline{b} can be used as test statistic, the critical region being given by

$$\{b : b > \chi^2_{2;\alpha}\}$$

For large n , this test has a size of approximately α .

A problem with this test is the very slow rate of convergence of $(\underline{b}_1, \underline{b}_2)$ to the bi-normal limiting distribution of Section 1. So, for moderate sample sizes this test is rather inaccurate. Especially the distribution of \underline{b}_2 is highly nonnormal, even if the sampled distribution is normal and even for large sample sizes. BOWMAN & SHENTON (1975) refined the test and obtained test contours for moderate n , based on a better approximation of the simultaneous distribution of $(\underline{b}_1, \underline{b}_2)$.

An alternative test may be based on $(\underline{s}, \underline{t})$. For large n , Section 2 implies

$$\underline{s} \approx N(0, 1.839), \quad \underline{t} \approx N(1.233, 3.153), \quad \underline{s} \text{ and } \underline{t} \text{ independent}$$

under H_0 and, consequently,

$$\underline{u} = n[\underline{s}^2/1.839 + (\underline{t} - 1.233)^2/3.153] \approx \chi_2^2 \quad (5.2)$$

The critical region for test statistic \underline{u} equals

$$\{u : u > \chi_{2;\alpha}^2\}$$

For large n , this test has a size of approximately α as well.

Since \underline{t} is much less influenced by outliers in the sample, it is to be expected that the convergence in distribution to the normal is much faster. This means that the approximation (5.2) is (much) better than (5.1) for moderate sample sizes. Therefore, this alternative quantile-based test may be a serious competitor to the existing ones. Approximate power-calculations can be made with the help of the general limiting distributions in Section 2. This topic too will be further investigated in the (near) future.

Note that there is nothing special in the null-hypothesis of normality: for **any** distribution within the Pearson system a similar test can be derived.

6 Discussion

This paper concentrated on measures for skewness and kurtosis only. Of course, limiting distributions for the quartets $(\underline{x}, \underline{s}^2, \underline{b}_1, \underline{b}_2)$ and $(\underline{q}, \underline{r}, \underline{s}, \underline{t})$ can be derived in a similar fashion.

In Section 4 attention was mainly directed towards Pearson's system of distributions. Similar results can be derived for Johnson's system, since there too all distributions have distinct pairs of values (β_1, β_2) and (S, T) . Still other systems may be considered as classes of possible models; interesting candidates are the SCHMEISER-DEUTSCH (1978) system and Burr's system, c.f. STUART & ORD (1987), p. 242.

Apart from the methods discussed in this paper - modelling by equating measures for location, dispersion, skewness and kurtosis - quite different methods of modelling can be used. We mention two possibilities: maximum likelihood and kernel estimation.

(i) Maximum likelihood (ML) estimation is usually applied when the type of density is given and only some parameters are unknown. ML-estimators have, under rather mild

conditions, nice asymptotic properties, like consistency and efficiency. In our case, a number of different types of densities are taken into consideration simultaneously. Since, however, Pearson's system is based upon a single differential equation with a limited number of parameters, ML-estimation may be possible in this case.

(ii) kernel estimation comes down to smoothing the empirical frequency distribution. From the sample $(\underline{x}_1, \underline{x}_2, \dots, \underline{x}_n)$, the density f is estimated by \underline{f} , where

$$\underline{f}(x) = \frac{1}{nh} \sum_{i=1}^n K[(x - \underline{x}_i)/h]$$

for all x . The kernel function K is concentrated around 0 and satisfies

$$\int_{-\infty}^{\infty} K(x)dx = 1$$

The bandwidth h determines the level of smoothing. Note that the resulting estimated density can not be described by a simple analytical formula.

References

- BOWMAN, K.O. and L.R. SHENTON (1975), *Omnibus test contours for departures from normality based on $\sqrt{b_1}$ and b_2* , Biometrika 62, 243-249.
- COENEN, V.M.J. (1992), *Fitting theoretical distributions to frequency distributions, based on moments and quantiles, respectively*, Tilburg University (in Dutch).
- D'AGOSTINO (1973), *Tests for departures from normality; empirical results for the distribution of b_2 and $\sqrt{b_1}$* , Biometrika 60, 613-622.
- MOORS, J.J.A. (1988), *A quantile alternative for kurtosis*, The Statistician 37, 25-32.
- SCHMEISER, B.W. and S.J. DEUTSCH (1987), *A versatile four parameter family of probability distributions suitable for simulation*, AIEE Transactions 9, 176-181.
- SERFLING, R.J. (1980), *Approximation theorems of mathematical statistics*, Wiley: New York.

- SHAPIRO, S.S., M.B. WILK and mrs. H.J. CHEN (1968), *A comparative study of various tests for normality*, Journal of the American Statistical Association 63, 1343-1372.
- STUART, A. and J.K. ORD (1987), *Kendall's advanced theory of statistics*, Griffin: London.
- WAGEMAKERS, R.Th.A., J.J.A. MOORS and M.J.B.T. JANSSENS (1992), *Characterizing distributions by quantile measures*, Report FEW 578, Tilburg University.

IN 1992 REEDS VERSCHENEN

- 532 F.G. van den Heuvel en M.R.M. Turlings
Privatisering van arbeidsongeschiktheidsregelingen
Refereed by Prof.Dr. H. Verbon
- 533 J.C. Engwerda, L.G. van Willigenburg
LQ-control of sampled continuous-time systems
Refereed by Prof.dr. J.M. Schumacher
- 534 J.C. Engwerda, A.C.M. Ran & A.L. Rijkeboer
Necessary and sufficient conditions for the existence of a positive definite solution
of the matrix equation $X + A^* X A = Q$.
Refereed by Prof.dr. J.M. Schumacher
- 535 Jacob C. Engwerda
The indefinite LQ-problem: the finite planning horizon case
Refereed by Prof.dr. J.M. Schumacher
- 536 Gert-Jan Otten, Peter Borm, Ton Storcken, Stef Tijs
Effectivity functions and associated claim game correspondences
Refereed by Prof.dr. P.H.M. Ruys
- 537 Jack P.C. Kleijnen, Gustav A. Alink
Validation of simulation models: mine-hunting case-study
Refereed by Prof.dr.ir. C.A.T. Takkenberg
- 538 V. Feltkamp and A. van den Nouweland
Controlled Communication Networks
Refereed by Prof.dr. S.H. Tijs
- 539 A. van Schaik
Productivity, Labour Force Participation and the Solow Growth Model
Refereed by Prof.dr. Th.C.M.J. van de Klundert
- 540 J.J.G. Lemmen and S.C.W. Eijffinger
The Degree of Financial Integration in the European Community
Refereed by Prof.dr. A.B.T.M. van Schaik
- 541 J. Bell, P.K. Jagersma
Internationale Joint Ventures
Refereed by Prof.dr. H.G. Barkema
- 542 Jack P.C. Kleijnen
Verification and validation of simulation models
Refereed by Prof.dr.ir. C.A.T. Takkenberg
- 543 Gert Nieuwenhuis
Uniform Approximations of the Stationary and Palm Distributions of Marked Point
Processes
Refereed by Prof.dr. B.B. van der Genugten

- 544 R. Heuts, P. Nederstigt, W. Roebroek, W. Selen
Multi-Product Cycling with Packaging in the Process Industry
Refereed by Prof.dr. F.A. van der Duyn Schouten
- 545 J.C. Engwerda
Calculation of an approximate solution of the infinite time-varying LQ-problem
Refereed by Prof.dr. J.M. Schumacher
- 546 Raymond H.J.M. Gradus and Peter M. Kort
On time-inconsistency and pollution control: a macroeconomic approach
Refereed by Prof.dr. A.J. de Zeeuw
- 547 Drs. Dolph Cantrijn en Dr. Rezaul Kabir
De Invloed van de Invoering van Preferente Beschermingsaandelen op Aandelen-
koersen van Nederlandse Beursgenoteerde Ondernemingen
Refereed by Prof.dr. P.W. Moerland
- 548 Sylvester Eijffinger and Eric Schaling
Central bank independence: criteria and indices
Refereed by Prof.dr. J.J. Sijben
- 549 Drs. A. Schmeits
Geïntegreerde investerings- en financieringsbeslissingen; Implicaties voor Capital
Budgeting
Refereed by Prof.dr. P.W. Moerland
- 550 Peter M. Kort
Standards versus standards: the effects of different pollution restrictions on the
firm's dynamic investment policy
Refereed by Prof.dr. F.A. van der Duyn Schouten
- 551 Niels G. Noorderhaven, Bart Nooteboom and Johannes Berger
Temporal, cognitive and behavioral dimensions of transaction costs; to an under-
standing of hybrid vertical inter-firm relations
Refereed by Prof.dr. S.W. Douma
- 552 Ton Storcken and Harrie de Swart
Towards an axiomatization of orderings
Refereed by Prof.dr. P.H.M. Ruys
- 553 J.H.J. Roemen
The derivation of a long term milk supply model from an optimization model
Refereed by Prof.dr. F.A. van der Duyn Schouten
- 554 Geert J. Almekinders and Sylvester C.W. Eijffinger
Daily Bundesbank and Federal Reserve Intervention and the Conditional Variance
Tale in DM/\$-Returns
Refereed by Prof.dr. A.B.T.M. van Schaik

- 555 Dr. M. Hetebrij, Drs. B.F.L. Jonker, Prof.dr. W.H.J. de Freytas
 "Tussen achterstand en voorsprong" de scholings- en personeelsvoorzieningsproblematiek van bedrijven in de procesindustrie
 Refereed by Prof.dr. Th.M.M. Verhallen
- 556 Ton Geerts
 Regularity and singularity in linear-quadratic control subject to implicit continuous-time systems
 Communicated by Prof.dr. J. Schumacher
- 557 Ton Geerts
 Invariant subspaces and invertibility properties for singular systems: the general case
 Communicated by Prof.dr. J. Schumacher
- 558 Ton Geerts
 Solvability conditions, consistency and weak consistency for linear differential-algebraic equations and time-invariant singular systems: the general case
 Communicated by Prof.dr. J. Schumacher
- 559 C. Fricker and M.R. Jaïbi
 Monotonicity and stability of periodic polling models
 Communicated by Prof.dr.ir. O.J. Boxma
- 560 Ton Geerts
 Free end-point linear-quadratic control subject to implicit continuous-time systems: necessary and sufficient conditions for solvability
 Communicated by Prof.dr. J. Schumacher
- 561 Paul G.H. Mulder and Anton L. Hempenius
 Expected Utility of Life Time in the Presence of a Chronic Noncommunicable Disease State
 Communicated by Prof.dr. B.B. van der Genugten
- 562 Jan van der Leeuw
 The covariance matrix of ARMA-errors in closed form
 Communicated by Dr. H.H. Tigelaar
- 563 J.P.C. Blanc and R.D. van der Mei
 Optimization of polling systems with Bernoulli schedules
 Communicated by Prof.dr.ir. O.J. Boxma
- 564 B.B. van der Genugten
 Density of the least squares estimator in the multivariate linear model with arbitrarily normal variables
 Communicated by Prof.dr. M.H.C. Paardekooper
- 565 René van den Brink, Robert P. Gilles
 Measuring Domination in Directed Graphs
 Communicated by Prof.dr. P.H.M. Ruys

- 566 Harry G. Barkema
The significance of work incentives from bonuses: some new evidence
Communicated by Dr. Th.E. Nijman
- 567 Rob de Groof and Martin van Tuijl
Commercial integration and fiscal policy in interdependent, financially integrated two-sector economies with real and nominal wage rigidity.
Communicated by Prof.dr. A.L. Bovenberg
- 568 F.A. van der Duyn Schouten, M.J.G. van Eijs, R.M.J. Heuts
The value of information in a fixed order quantity inventory system
Communicated by Prof.dr. A.J.J. Talman
- 569 E.N. Kertzman
Begrotingsnormering en EMU
Communicated by Prof.dr. J.W. van der Dussen
- 570 A. van den Elzen, D. Talman
Finding a Nash-equilibrium in noncooperative N-person games by solving a sequence of linear stationary point problems
Communicated by Prof.dr. S.H. Tijs
- 571 Jack P.C. Kleijnen
Verification and validation of models
Communicated by Prof.dr. F.A. van der Duyn Schouten
- 572 Jack P.C. Kleijnen and Willem van Groenendaal
Two-stage versus sequential sample-size determination in regression analysis of simulation experiments
Communicated by Prof.Dr. F.A. van der Duyn Schouten
- 573 Pieter K. Jagersma
Het management van multinationale ondernemingen: de concernstructuur
Communicated by Prof.Dr. S.W. Douma
- 574 A.L. Hempenius
Explaining Changes in External Funds. Part One: Theory
Communicated by Prof.Dr.Ir. A. Kapteyn
- 575 J.P.C. Blanc, R.D. van der Mei
Optimization of Polling Systems by Means of Gradient Methods and the Power-Series Algorithm
Communicated by Prof.dr.ir. O.J. Boxma
- 576 Herbert Hamers
A silent duel over a cake
Communicated by Prof.dr. S.H. Tijs

- 577 Gerard van der Laan, Dolf Talman, Hans Kremers
On the existence and computation of an equilibrium in an economy with constant returns to scale production
Communicated by Prof.dr. P.H.M. Ruys
- 578 R.Th.A. Wagemakers, J.J.A. Moors, M.J.B.T. Janssens
Characterizing distributions by quantile measures
Communicated by Dr. R.M.J. Heuts
- 579 J. Ashayeri, W.H.L. van Esch, R.M.J. Heuts
Amendment of Heuts-Selen's Lotsizing and Sequencing Heuristic for Single Stage Process Manufacturing Systems
Communicated by Prof.dr. F.A. van der Duyn Schouten
- 580 H.G. Barkema
The Impact of Top Management Compensation Structure on Strategy
Communicated by Prof.dr. S.W. Douma
- 581 Jos Benders en Freek Aertsen
Aan de lijn of aan het lijntje: wordt slank produceren de mode?
Communicated by Prof.dr. S.W. Douma
- 582 Willem Haemers
Distance Regularity and the Spectrum of Graphs
Communicated by Prof.dr. M.H.C. Paardekooper
- 583 Jalal Ashayeri, Behnam Pourbabai, Luk van Wassenhove
Strategic Marketing, Production, and Distribution Planning of an Integrated Manufacturing System
Communicated by Prof.dr. F.A. van der Duyn Schouten
- 584 J. Ashayeri, F.H.P. Driessen
Integration of Demand Management and Production Planning in a Batch Process Manufacturing System: Case Study
Communicated by Prof.dr. F.A. van der Duyn Schouten
- 585 J. Ashayeri, A.G.M. van Eijs, P. Nederstigt
Blending Modelling in a Process Manufacturing System
Communicated by Prof.dr. F.A. van der Duyn Schouten
- 586 J. Ashayeri, A.J. Westerhof, P.H.E.L. van Alst
Application of Mixed Integer Programming to A Large Scale Logistics Problem
Communicated by Prof.dr. F.A. van der Duyn Schouten
- 587 P. Jean-Jacques Herings
On the Structure of Constrained Equilibria
Communicated by Prof.dr. A.J.J. Talman

IN 1993 REEDS VERSCHENEN

- 588 Rob de Groof and Martin van Tuijl
The Twin-Debt Problem in an Interdependent World
Communicated by Prof.dr. Th. van de Klundert
- 589 Harry H. Tigelaar
A useful fourth moment matrix of a random vector
Communicated by Prof.dr. B.B. van der Genugten
- 590 Niels G. Noorderhaven
Trust and transactions; transaction cost analysis with a differential behavioral assumption
Communicated by Prof.dr. S.W. Douma
- 591 Henk Roest and Kitty Koelemeijer
Framing perceived service quality and related constructs A multilevel approach
Communicated by Prof.dr. Th.M.M. Verhallen
- 592 Jacob C. Engwerda
The Square Indefinite LQ-Problem: Existence of a Unique Solution
Communicated by Prof.dr. J. Schumacher
- 593 Jacob C. Engwerda
Output Deadbeat Control of Discrete-Time Multivariable Systems
Communicated by Prof.dr. J. Schumacher
- 594 Chris Veld and Adri Verboven
An Empirical Analysis of Warrant Prices versus Long Term Call Option Prices
Communicated by Prof.dr. P.W. Moerland
- 595 A.A. Jeunink en M.R. Kabir
De relatie tussen aandelhoudersstructuur en beschermingsconstructies
Communicated by Prof.dr. P.W. Moerland
- 596 M.J. Coster and W.H. Haemers
Quasi-symmetric designs related to the triangular graph
Communicated by Prof.dr. M.H.C. Paardekooper
- 597 Noud Gruijters
De liberalisering van het internationale kapitaalverkeer in historisch-institutioneel perspectief
Communicated by Dr. H.G. van Gemert
- 598 John Görtzen en Remco Zwetheul
Weekend-effect en dag-van-de-week-effect op de Amsterdamse effectenbeurs?
Communicated by Prof.dr. P.W. Moerland
- 599 Philip Hans Franses and H. Peter Boswijk
Temporal aggregation in a periodically integrated autoregressive process
Communicated by Prof.dr. Th.E. Nijman

- 600 René Peeters
On the p-ranks of Latin Square Graphs
Communicated by Prof.dr. M.H.C. Paardekooper
- 601 Peter E.M. Borm, Ricardo Cao, Ignacio García-Jurado
Maximum Likelihood Equilibria of Random Games
Communicated by Prof.dr. B.B. van der Genugten
- 602 Prof.dr. Robert Bannink
Size and timing of profits for insurance companies. Cost assignment for products with multiple deliveries.
Communicated by Prof.dr. W. van Hulst
- 603 M.J. Coster
An Algorithm on Addition Chains with Restricted Memory
Communicated by Prof.dr. M.H.C. Paardekooper
- 604 Ton Geerts
Coordinate-free interpretations of the optimal costs for LQ-problems subject to implicit systems
Communicated by Prof.dr. J.M. Schumacher
- 605 B.B. van der Genugten
Beat the Dealer in Holland Casino's Black Jack
Communicated by Dr. P.E.M. Borm
- 606 Gert Nieuwenhuis
Uniform Limit Theorems for Marked Point Processes
Communicated by Dr. M.R. Jaïbi
- 607 Dr. G.P.L. van Roij
Effectisering op internationale financiële markten en enkele gevolgen voor banken
Communicated by Prof.dr. J. Sijben
- 608 R.A.M.G. Joosten, A.J.J. Talman
A simplicial variable dimension restart algorithm to find economic equilibria on the unit simplex using $n(n+1)$ rays
Communicated by Prof.Dr. P.H.M. Ruys
- 609 Dr. A.J.W. van de Gevel
The Elimination of Technical Barriers to Trade in the European Community
Communicated by Prof.dr. H. Huizinga
- 610 Dr. A.J.W. van de Gevel
Effective Protection: a Survey
Communicated by Prof.dr. H. Huizinga
- 611 Jan van der Leeuw
First order conditions for the maximum likelihood estimation of an exact ARMA model
Communicated by Prof.dr. B.B. van der Genugten

- 612 Tom P. Faith
 Bertrand-Edgeworth Competition with Sequential Capacity Choice
 Communicated by Prof.Dr. S.W. Douma
- 613 Ton Geerts
 The algebraic Riccati equation and singular optimal control: The discrete-time case
 Communicated by Prof.dr. J.M. Schumacher
- 614 Ton Geerts
 Output consistency and weak output consistency for continuous-time implicit systems
 Communicated by Prof.dr. J.M. Schumacher
- 615 Stef Tijs, Gert-Jan Otten
 Compromise Values in Cooperative Game Theory
 Communicated by Dr. P.E.M. Borm
- 616 Dr. Pieter J.F.G. Meulendijks and Prof.Dr. Dick B.J. Schouten
 Exchange Rates and the European Business Cycle: an application of a 'quasi-empirical' two-country model
 Communicated by Prof.Dr. A.H.J.J. Kolnaar
- 617 Niels G. Noorderhaven
 The argumentational texture of transaction cost economics
 Communicated by Prof.Dr. S.W. Douma
- 618 Dr. M.R. Jaïbi
 Frequent Sampling in Discrete Choice
 Communicated by Dr. M.H. ten Raa
- 619 Dr. M.R. Jaïbi
 A Qualification of the Dependence in the Generalized Extreme Value Choice Model
 Communicated by Dr. M.H. ten Raa

Bibliotheek K. U. Brabant



17 000 01176232 6